



Dr. Joshua Alpert
 Post-Operative Rehabilitation Protocol:
 Shoulder - Reverse Shoulder Replacement

	Range of Motion	Sling	Exercises	Precautions
PHASE I 0-6 weeks No physical therapy.	None. No moving at the shoulder joint.	On at all times except for hygiene	Pendulum exercises at home, gentle elbow and wrist ROM at home.	- no lifting at the shoulder joint - no weight bearing on the hand/wrist/elbow to avoid dislocation. No pushing up out of a chair, no using a walker etc.
PHASE II 6-12 weeks Physical therapy starts.	As tolerated	None	AROM and PROM of shoulder. No strengthening.	- no weight bearing on the hand/wrist/elbow to avoid dislocation. No pushing up out of a chair, no using a walker etc.
PHASE III 12-18 weeks	Progress to Full	None	Start strengthening of the shoulder as tolerated. Closed chain scapular rehab and functional rotator cuff	- no weight bearing on the hand/wrist/elbow. No pushing up out of a chair, no using a walker etc.

			strengthening. Focus on anterior deltoid and teres. Maximize scapular stabilization.	
PHASE IV 18 weeks+	Full	None	Progress with strengthening and scapular stabilization.	- avoid weight bearing through the hand and exercises alike. Ex: pushups, using a walker, pushing out of a chair etc.

Fox Valley Orthopedics
420 W. Northwest Highway
Barrington, IL 60010
847.382.6766

Fox Valley Orthopedics
1710 Randall Road
Elgin, IL 60123
224.293.1170